

BEURLING DIMENSION OF GABOR PSEUDOFRAMES FOR AFFINE SUBSPACES

WOJCIECH CZAJA, GITTA KUTYNIOK, AND DARRIN SPEEGLE

ABSTRACT. Pseudoframes for subspaces have been recently introduced by S. Li and H. Ogawa as a tool to analyze lower dimensional data with arbitrary flexibility of both the analyzing and the dual sequence.

In this paper we study Gabor pseudoframes for affine subspaces by focusing on geometrical properties of their associated sets of parameters. We first introduce a new notion of Beurling dimension for discrete subsets of \mathbb{R}^d by employing a certain generalized Beurling density. We present several properties of Beurling dimension including a comparison with other notions of dimension showing, for instance, that our notion includes the mass dimension as a special case. Then we prove that Gabor pseudoframes for affine subspaces satisfy a certain Homogeneous Approximation Property, which implies invariance under time-frequency shifts of an approximation by elements from the pseudoframe.

The main result of this paper is a classification of Gabor pseudoframes for affine subspaces by means of the Beurling dimension of their sets of parameters. This provides us, in particular, with a Nyquist dimension which separates sets of parameters of pseudoframes from those of non-pseudoframes and which links a fixed value to sets of parameters of pseudo-Riesz sequences. These results are even new for the special case of Gabor frames for an affine subspace.

1. INTRODUCTION

Frames are a well-established tool both in applied and pure mathematics which provides robust and stable – but usually nonunique – representations of vectors. For instance, in wireless communications frames ensure robustness of transmission against erasures [21], in image processing they serve as building blocks for novel directional representation systems [3], and just recently it has been discovered that the theory of frames may provide a means to attack the Kadison–Singer Conjecture in operator theory from 1959 [4].

However, for some applications frames lack enough flexibility, for instance, in the design of the dual frame needed for reconstruction. One approach to circumvent this problem was the introduction of *pseudoframes* [15]. In the situation of a pseudoframe,

⁰*Math Subject Classifications.* 42C15, 42C40

⁰*Keywords and Phrases.* Beurling density, Beurling dimension, frame, Gabor system, discrete Hausdorff dimension, Homogeneous Approximation Property, mass dimensions, Nyquist density, pseudoframe, pseudoframe for subspaces

Date: October 20, 2006.

the “dual sequence” is only required to provide a reconstruction formula, thereby allowing more flexibility. In some instances, data that we wish to analyze is contained in a subspace, which naturally leads to the idea of *pseudoframes for subspaces* [16].

In this paper we will study a special class of pseudoframes for subspaces which are of particular interest in time-frequency analysis. A *Gabor system* consists of a collection of time-frequency shifts of a single function or a finite family of functions in $L^2(\mathbb{R}^d)$ with respect to a discrete set of parameters in $\mathbb{R}^d \times \mathbb{R}^d$. Due to this structure, Gabor systems are especially suitable for applications involving time-dependent frequency content, for example, for the analysis of acoustic signals like music [7].

Classical Gabor systems, which employ just a lattice as set of parameters, have been studied extensively over the past 20 years. Recently, the more general, *irregular Gabor systems* with arbitrary sets of parameters in $\mathbb{R}^d \times \mathbb{R}^d$ have attracted increasing attention, in particular due to applications from sampling and perturbation theory. Questions concerning (frame) properties of irregular Gabor systems lead naturally to the study of the associated sets of parameters. A very elegant way to classify sets of parameters is the consideration of their Beurling densities. For a recent survey article with extensive list of references we refer to [9].

However, we intend to focus on the study of *Gabor pseudoframes for (affine) subspaces*. Beurling density alone does not serve our needs here, since the “dimensionality” of the set of parameters will play an essential role. Beurling density on the contrary only serves as a classifying tool for sets of parameters of the same “dimension”. Comparable problems occur in the study of sets of parameters of *wave packet systems*, which are systems consisting of dilates, translates, and modulates of a single function in $L^2(\mathbb{R})$ with their sets of parameters being contained in the affine Weyl–Heisenberg group. Progress on the problem of classifying sets of parameters of wave packet systems was recently made by the authors through the introduction of *upper and lower Beurling dimensions* based on Beurling density and inspired by the notion of Hausdorff dimension [6].

In this paper, we present a notion of Beurling dimension for discrete subsets of \mathbb{R}^d by employing a certain generalized Beurling density. First, we study several of its properties, including behavior under perturbations, monotonicity, stability, and geometric invariance. We further compare the new notion with the most well-known notions of dimension for discrete subsets of \mathbb{R}^d , such as, for example, the mass dimension, which we show to be a special case of our notion. Secondly, we apply the new notion of Beurling dimension to Gabor pseudoframes for (affine) subspaces. In particular, inspired by techniques from [10, 20], we determine the Beurling dimensions of the sets of parameters of Gabor pseudoframes for affine subspaces. This leads to a classification of Gabor pseudoframes for affine subspaces and Gabor pseudo-Riesz sequences by means of Beurling dimensions of their associated sets of parameters (Theorem 3.3).

This main results and the techniques of their proofs have several interesting implications. One of them is that Gabor pseudoframes for affine subspaces cannot be expected to have a Nyquist density: a density separating sets of parameters of frames from those of non-frames, such as in the situation of Gabor frames (cf. [5]). This

is due to the fact that the notion of density does not provide useful information in this context. However, our result shows that instead we have a *Nyquist dimension*, which separates sets of parameters of pseudoframes from those of non-pseudoframes and links a fixed value to sets of parameters of pseudo-Riesz sequences. This result is new even in the situation of Gabor frames for an affine subspace. Also, along the way, we prove that Gabor pseudoframes for affine subspaces always satisfy a certain *Homogeneous Approximation Property*, which can be interpreted as invariance under time-frequency shifts of an approximation by elements from a Gabor pseudoframe for an affine subspace.

This paper is organized as follows. The definition and general theory of Beurling dimension for discrete subsets of \mathbb{R}^d is given in Section 2, including a comparison of Beurling dimension with mass dimensions and the discrete Hausdorff dimension. In Section 3 we determine the Beurling dimensions of the sets of parameters of Gabor pseudoframes for affine subspaces and prove the classification result.

2. BEURLING DIMENSION

2.1. Definition of Beurling dimension. We will define a notion of Beurling dimension for sequences Λ in \mathbb{R}^d suited to Euclidean geometry, by using as a backbone a generalization of Beurling density. Notice that throughout this paper, although Λ will always denote a sequence of points in \mathbb{R}^d and not merely a subset, for simplicity we will write $\Lambda \subset \mathbb{R}^d$.

First we require some notation. Let Q denote the box $[-1, 1]^d$ and, for $h > 0$, let Q_h be the dilation of Q by the factor of h :

$$Q_h = hQ = [-h, h]^d.$$

For any $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ we let $Q_h(x)$ be the set Q_h translated in such a way so that it is “centered” at x , i.e.,

$$Q_h(x) = \prod_{i=1}^d [x_i - h, x_i + h].$$

Employing these notions we define a generalization of Beurling density.

Definition 2.1. Let $\Lambda \subset \mathbb{R}^d$, S be an affine subspace of \mathbb{R}^d and $r > 0$. The lower Beurling density of Λ with respect to S and r is defined by

$$\mathcal{D}_{S,r}^-(\Lambda) = \liminf_{h \rightarrow \infty} \inf_{x \in S} \frac{\#(\Lambda \cap Q_h(x))}{h^r},$$

and the upper Beurling density of Λ with respect to S and r is defined by

$$\mathcal{D}_{S,r}^+(\Lambda) = \limsup_{h \rightarrow \infty} \sup_{x \in S} \frac{\#(\Lambda \cap Q_h(x))}{h^r}.$$

If $\mathcal{D}_{S,r}^-(\Lambda) = \mathcal{D}_{S,r}^+(\Lambda)$, then we say that Λ has uniform Beurling density with respect to S and r , and we denote this density by $\mathcal{D}_{S,r}(\Lambda)$.

We remark that the Beurling density with respect to \mathbb{R}^d and d coincides to the classical Beurling density. In particular, for $S = \mathbb{R}^d$ and $r = d$ the definition of Beurling density is independent of the particular choice of the set Q (see [14]). By using a similar argument for each $r > 0$, we obtain:

Proposition 2.2. *Let $\Lambda \subset \mathbb{R}^d$, and let $U \subset \mathbb{R}^d$ be a compact set of measure 2^d whose boundary has measure zero. Then, for all $r > 0$ and all affine subspaces S of \mathbb{R}^d ,*

$$\mathcal{D}_{S,r}^-(\Lambda) = \liminf_{h \rightarrow \infty} \inf_{x \in S} \frac{\#(\Lambda \cap (x + hU))}{h^r}$$

and

$$\mathcal{D}_{S,r}^+(\Lambda) = \limsup_{h \rightarrow \infty} \sup_{x \in S} \frac{\#(\Lambda \cap (x + hU))}{h^r}.$$

For $y \in \mathbb{R}^d$, Beurling density with respect to $S = \{y\}$ and $r = d$ is also a commonly used density, and it is widely known that such density does not depend on y . More generally, we have the following:

Proposition 2.3. *Let $\Lambda \subset \mathbb{R}^d$, T a linear subspace of \mathbb{R}^d , and $y \in \mathbb{R}^d$. Then, for all $r > 0$,*

$$\mathcal{D}_{T+y,r}^+(\Lambda) = \mathcal{D}_{T,r}^+(\Lambda)$$

and

$$\mathcal{D}_{T+y,r}^-(\Lambda) = \mathcal{D}_{T,r}^-(\Lambda).$$

Proof. Note that there exists $R > 0$ such that for each $x \in T$ and $h > 0$, $Q_{h+R}(x) \supset Q_h(x+y)$ and $Q_{h+R}(x+y) \supset Q_h(x)$. The result follows. \square

We now define the Beurling dimension of a set, which will be used to characterize the Nyquist dimension of Gabor pseudoframes for an affine subspace.

Definition 2.4. *Let $\Lambda \subset \mathbb{R}^d$ and let S be an affine subspace of \mathbb{R}^d . The lower Beurling dimension of $\Lambda \subset \mathbb{R}^d$ with respect to S is defined by*

$$\dim_S^-(\Lambda) = \inf \{r > 0 : \mathcal{D}_{S,r}^-(\Lambda) < \infty\}$$

and the upper Beurling dimension of $\Lambda \subset \mathbb{R}^d$ with respect to S is

$$\dim_S^+(\Lambda) = \sup \{r > 0 : \mathcal{D}_{S,r}^+(\Lambda) > 0\}.$$

When these two quantities are equal, we refer to the Beurling dimension of Λ with respect to S , and we denote it by $\dim_S(\Lambda)$.

It follows immediately from the definition that we always have $\dim_S^-(\Lambda) \leq \dim_S^+(\Lambda)$.

The following result presents possible different definitions of Beurling dimensions. Since the proof is rather technical and uses the same arguments as the proof of a similar result in [6], we omit it here.

Proposition 2.5. *Let $\Lambda \subset \mathbb{R}^d$ and let S be an affine subspace of \mathbb{R}^d . Then,*

- (i) $\dim_S^-(\Lambda) = \sup \{r > 0 : \mathcal{D}_{S,r}^-(\Lambda) > 0\}$,
- (ii) $\dim_S^+(\Lambda) = \inf \{r > 0 : \mathcal{D}_{S,r}^+(\Lambda) < \infty\}$.

We note here that, while Beurling dimension is defined above for arbitrary subsets of \mathbb{R}^d , the upper Beurling dimension will be infinite unless Λ is discrete. Indeed, if x is an accumulation point of Λ , then for all $h > 0$, $\#(\Lambda \cap Q_h(x)) = \infty$. Thus, we will restrict our attention to discrete subsets of Λ .

2.2. Properties of Beurling dimension. In this section we present several properties of the Beurling dimension with respect to $S \subset \mathbb{R}^d$.

2.2.1. Perturbation. We first show that Beurling density is robust against perturbations of the elements in the set under consideration. Given $\epsilon > 0$, we say that $\Delta \subset \mathbb{R}^d$ is an ϵ -perturbation of $\Lambda \subset \mathbb{R}^d$, if $\Delta = \{\lambda + \delta_\lambda : \lambda \in \Lambda, \delta_\lambda \in [-\epsilon, \epsilon]\}$. As before, Δ should be considered as a sequence rather than a subset of \mathbb{R}^d .

Lemma 2.6. *Let $\Lambda \subset \mathbb{R}^d$, S an affine subspace of \mathbb{R}^d , and $\epsilon > 0$. For any ϵ -perturbation Δ of Λ and any $r > 0$, we have*

$$\mathcal{D}_{S,r}^-(\Lambda) = \mathcal{D}_{S,r}^-(\Delta) \quad \text{and} \quad \mathcal{D}_{S,r}^+(\Lambda) = \mathcal{D}_{S,r}^+(\Delta).$$

Proof. For $h > 0$, $x \in S$, we obtain the following estimates for $\#(\Lambda \cap Q_h(x))$:

$$\#(\Delta \cap Q_{h-\epsilon}(x)) \leq \#(\Lambda \cap Q_h(x)) \leq \#(\Delta \cap Q_{h+\epsilon}(x)).$$

Dividing the terms by h^r for $r > 0$ and observing that

$$\limsup_{h \rightarrow \infty} \frac{\sup_{x \in S} \#(\Delta \cap Q_{h-\epsilon}(x))}{h^r} = \mathcal{D}_r^+(\Delta) = \limsup_{h \rightarrow \infty} \frac{\sup_{x \in S} \#(\Delta \cap Q_{h+\epsilon}(x))}{h^r},$$

implies $\mathcal{D}_{S,r}^+(\Lambda) = \mathcal{D}_{S,r}^+(\Delta)$.

The claim concerning the lower Beurling density with respect to S and r can be treated similarly. \square

This implies the following perturbation result.

Theorem 2.7. *Let $\Lambda \subset \mathbb{R}^d$, S an affine subspace of \mathbb{R}^d , and $\epsilon > 0$. For any ϵ -perturbation Δ of Λ , we have*

$$\dim_S^-(\Lambda) = \dim_S^-(\Delta) \quad \text{and} \quad \dim_S^+(\Lambda) = \dim_S^+(\Delta).$$

Proof. This follows from Lemma 2.6 and the definition of upper and lower Beurling dimension. \square

2.2.2. Geometric Properties. Our next result shows that the upper, lower, and uniform Beurling dimensions satisfy the following properties which are typically attributed to dimensions.

Proposition 2.8. *Let $\Lambda_1, \Lambda_2, \Lambda \subset \mathbb{R}^d$ and S an affine subspace of \mathbb{R}^d . Then the following conditions hold.*

(i) **Monotonicity:** *If $\Lambda_1 \subseteq \Lambda_2$, we have*

$$\dim_S^-(\Lambda_1) \leq \dim_S^-(\Lambda_2) \quad \text{and} \quad \dim_S^+(\Lambda_1) \leq \dim_S^+(\Lambda_2).$$

(ii) **Stability:** *We have*

$$\dim_{\mathcal{S}}^{-}(\Lambda_1 \cap \Lambda_2) \leq \min(\dim_{\mathcal{S}}^{-}(\Lambda_1), \dim_{\mathcal{S}}^{-}(\Lambda_2))$$

and

$$\dim_{\mathcal{S}}^{+}(\Lambda_1 \cup \Lambda_2) = \max(\dim_{\mathcal{S}}^{+}(\Lambda_1), \dim_{\mathcal{S}}^{+}(\Lambda_2)).$$

(iii) **Geometric invariance:** *Let $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a uniform homeomorphism such that $f(S) = S + y$ for some $y \in \mathbb{R}^d$. Then*

$$\dim_{\mathcal{S}}^{-}(f(\Lambda)) = \dim_{\mathcal{S}}^{-}(\Lambda) \quad \text{and} \quad \dim_{\mathcal{S}}^{+}(f(\Lambda)) = \dim_{\mathcal{S}}^{+}(\Lambda).$$

Proof. We will only prove the claims for the upper Beurling dimension. The lower Beurling dimension can be treated similarly.

(i) If $\Lambda_1 \subseteq \Lambda_2$, it follows that $\mathcal{D}_{S,r}^{+}(\Lambda_1) \leq \mathcal{D}_{S,r}^{+}(\Lambda_2)$ for all $r > 0$. Using the definition of Beurling dimension proves the claim.

(ii) For all $r > 0$, we have $\mathcal{D}_{S,r}^{+}(\Lambda_1 \cup \Lambda_2) \geq \max(\mathcal{D}_{S,r}^{+}(\Lambda_1), \mathcal{D}_{S,r}^{+}(\Lambda_2))$. Thus

$$(1) \quad \dim_{\mathcal{S}}^{+}(\Lambda_1 \cup \Lambda_2) \geq \max(\dim_{\mathcal{S}}^{+}(\Lambda_1), \dim_{\mathcal{S}}^{+}(\Lambda_2)).$$

Now fix $r > 0$. Since

$$\sup_{x \in S} \#((\Lambda_1 \cup \Lambda_2) \cap Q_h(x)) \leq \sup_{x \in S} \#(\Lambda_1 \cap Q_h(x)) + \sup_{x \in S} \#(\Lambda_2 \cap Q_h(x)),$$

it follows that $\mathcal{D}_{S,r}^{+}(\Lambda_1 \cup \Lambda_2) \leq \mathcal{D}_{S,r}^{+}(\Lambda_1) + \mathcal{D}_{S,r}^{+}(\Lambda_2)$. This implies $\mathcal{D}_{S,r}^{+}(\Lambda_1 \cup \Lambda_2) \leq 2 \max(\mathcal{D}_{S,r}^{+}(\Lambda_1), \mathcal{D}_{S,r}^{+}(\Lambda_2))$. Due to the definition of $\dim_{\mathcal{S}}^{+}$, this yields

$$(2) \quad \dim_{\mathcal{S}}^{+}(\Lambda_1 \cup \Lambda_2) \leq \max(\dim_{\mathcal{S}}^{+}(\Lambda_1), \dim_{\mathcal{S}}^{+}(\Lambda_2)).$$

Equations (1) and (2) settle the claim.

(iii) Note that, if f is uniformly continuous between two normed linear spaces and onto, then it is Lipschitz for large distances, see for example [2, Lemma 5.1]. That is, for all $h_0 > 0$, there exists $C > 0$ such that $f(Q_h(x)) \subset Q_{Ch}(f(x))$ for all $x \in \mathbb{R}^d$ and $h \geq h_0$. In particular, we conclude that for each $r > 0$,

$$\begin{aligned} \limsup_{h \rightarrow \infty} \sup_{x \in S} \frac{\#(f^{-1}(\Lambda) \cap Q_h(x))}{h^r} &\leq \limsup_{h \rightarrow \infty} \sup_{x \in S} \frac{\#(\Lambda \cap Q_{Ch}(f(x)))}{h^r} \\ &\leq C^r \limsup_{h \rightarrow \infty} \sup_{x \in S+y} \frac{\#(\Lambda \cap Q_h(x))}{h^r}. \end{aligned}$$

It follows from Proposition 2.3 that $\dim_{\mathcal{S}}^{+}(f^{-1}(\Lambda)) \leq \dim_{\mathcal{S}}^{+}(\Lambda)$. A similar argument shows that $\dim_{\mathcal{S}}^{+}(f(f^{-1}(\Lambda))) \leq \dim_{\mathcal{S}}^{+}(f^{-1}(\Lambda))$, which yields the result. \square

2.2.3. *Range of values.* Beurling dimension is particularly well-behaved in the case $S = \mathbb{R}^d$, or more generally under the hypotheses of Proposition 2.9 below. With this additional hypothesis, we present the possible range of values of Beurling dimension, making use of Proposition 2.9 below. This result provides a variety of different interpretations of finite upper and positive lower Beurling density with respect to $r = d$. Recall that $\Lambda \subset \mathbb{R}^d$ is *uniformly separated*, if $\inf_{\lambda_1, \lambda_2 \in \Lambda, \lambda_1 \neq \lambda_2} |\lambda_1 - \lambda_2| > 0$. It is *relatively uniformly separated*, if it is a finite union of uniformly separated sets. Further, Λ is *h-dense*, if $\bigcup_{x \in \Lambda} Q_h(x) = \mathbb{R}^d$.

Proposition 2.9. *Let S, V be affine subspaces of \mathbb{R}^d and suppose that $\Lambda \subset V$ and that there exists z_0 such that $V + z_0 \subset S$. Let n be the affine dimension of V .*

- (i) *The following conditions are equivalent.*
 - (a) $\mathcal{D}_{S,n}^+(\Lambda) < \infty$.
 - (b) $\mathcal{D}_{S,\dim(S)}^+(\Lambda) < \infty$.
 - (c) *There exists some $h > 0$ such that $\sup_{x \in S} \#(\Lambda \cap Q_h(x)) < \infty$.*
 - (d) *For all $h > 0$, $\sup_{x \in S} \#(\Lambda \cap Q_h(x)) < \infty$.*
 - (e) *Λ is relatively uniformly separated.*
 - (f) *For all $h > 0$, $\sup_{x \in S} \#\{\lambda \in \Lambda : x \in Q_h(\lambda)\} < \infty$.*
- (ii) *Also the following conditions are equivalent.*
 - (a) $\mathcal{D}_{S,n}^-(\Lambda) > 0$.
 - (b) $\mathcal{D}_{S,\dim(S)}^-(\Lambda) > 0$.
 - (c) *There exists some $h > 0$ such that $\inf_{x \in S} \#(\Lambda \cap Q_h(x)) > 0$.*
 - (d) *Λ contains a subsequence of positive uniform density.*
 - (e) *There exists some $h > 0$ such that Λ is h-dense.*

Proof. Note that under the hypotheses of this proposition, Propositions 2.2 and 2.3, and the proof of Proposition 2.8 (iii) imply that, without loss of generality, we may consider the case that $S = \mathbb{R}^d$ and $\Lambda \subset V = \mathbb{R}^n \times \{0, \dots, 0\} \subset \mathbb{R}^d$.

(i): (b), (c), (d), and (e) are equivalent by [5, Lemma 2.3]. To prove (d) \Leftrightarrow (f) observe that

$$\#(\Lambda \cap Q_h(x)) = \#\{\lambda \in \Lambda : -x \in Q_h - \lambda\} = \#\{\lambda \in \Lambda : x \in Q_h(\lambda)\}.$$

This immediately settles the claim.

The fact that (a) implies (b) is immediate. To show that (b) implies (a), we proceed *a contrario* and assume that $\mathcal{D}_{\mathbb{R}^d,n}^+(\Lambda) = \infty$. First, we note that, as a consequence of $\Lambda \subset V$, we may write $\Lambda = \Lambda' \times \{0, \dots, 0\} \subset \mathbb{R}^n \times \{0, \dots, 0\} = \mathbb{R}^d$ and so, $\mathcal{D}_{\mathbb{R}^d,n}^+(\Lambda) = \mathcal{D}_{\mathbb{R}^n,n}^+(\Lambda')$ and $\mathcal{D}_{\mathbb{R}^d,d}^+(\Lambda) = \mathcal{D}_{\mathbb{R}^n,d}^+(\Lambda')$. Next, we again invoke [5, Lemma 2.3] to conclude that $\sup_{x \in \mathbb{R}^n} \#(\Lambda' \cap Q_h(x)) = \infty$, which, in turn, implies that $\mathcal{D}_{\mathbb{R}^n,d}^+(\Lambda') = \infty$.

(ii): Clearly, (b) implies (a). On the other hand, since $\Lambda \subset \mathbb{R}^n \times \{0, \dots, 0\}$, $\mathcal{D}_{\mathbb{R}^d,n}^-(\Lambda) > 0$ only if $n = d$.

The implication (b) \Rightarrow (c) is immediate.

To show (c) \Rightarrow (d), let $h, \delta > 0$ be such that $\#(\Lambda \cap Q_h(x)) > \delta$ for all $x \in \mathbb{R}^d$. Hence, in particular, each set $\Lambda \cap Q_h(x)$ contains at least one element. Thus, for each $k \in \mathbb{Z}^d$

there exists some $y_k \in \Lambda \cap Q_h(2hk)$. Since $(y_k)_{k \in \mathbb{Z}^d}$ is a $2h$ -perturbation of $(2h\mathbb{Z})^d$ and $\mathcal{D}_{\mathbb{R}^d, d}((2h\mathbb{Z})^d) = (2h)^{-d}$, Lemma 2.6 implies that $(y_k)_{k \in \mathbb{Z}^d}$ has uniform density equal to $(2h)^{-d}$, which proves the claim.

Next suppose that Λ contains a subsequence Δ of positive uniform density. Since $\mathcal{D}_{\mathbb{R}^d, d}^-(\Lambda) > \mathcal{D}_{\mathbb{R}^d, d}(\Delta) > 0$, (d) implies (b).

The equivalence of (c) and (e) is immediate. \square

Theorem 2.10. *Let S, V be affine subspaces of \mathbb{R}^d and suppose that $\Lambda \subset V$ and that there exists z_0 such that $V + z_0 \subset S$. Denote the affine dimension of V by n . Then,*

- (i) $\dim_S^+(\Lambda) \in [0, n] \cup \{\infty\}$.
- (ii) $\dim_S^-(\Lambda) \in \{0\} \cup [n, \infty]$.

Proof. As before, we may assume without loss of generality that $S = \mathbb{R}^d$. Assume that $\dim_{\mathbb{R}^d}^+(\Lambda) =: s > n$ and $s < \infty$. By definition, this implies $\mathcal{D}_{\mathbb{R}^d, n}^+(\Lambda) = \infty$. By Proposition 2.9(i), $\sup_{x \in \mathbb{R}^d} \#(\Lambda \cap Q_h(x)) = \infty$ for all $h > 0$. Hence $\mathcal{D}_{\mathbb{R}^d, s+1}^+(\Lambda) = \infty$, a contradiction to $s = \sup\{r > 0 : \mathcal{D}_{\mathbb{R}^d, r}^+(\Lambda) > 0\}$ and $s < \infty$. This proves (i).

To show (ii) assume that $\dim_{\mathbb{R}^d}^-(\Lambda) =: s \in (0, n)$. This implies $\mathcal{D}_{\mathbb{R}^d, n}^-(\Lambda) = 0$. By Proposition 2.9(ii), $\inf_{x \in \mathbb{R}^d} \#(\Lambda \cap Q_h(x)) = 0$ for all $h > 0$. Hence also $\mathcal{D}_{\mathbb{R}^d, \frac{s}{2}}^-(\Lambda) = 0$, which contradicts $s = \inf\{r > 0 : \mathcal{D}_{\mathbb{R}^d, r}^-(\Lambda) < \infty\}$ and $s > 0$. \square

This theorem implies that the upper Beurling dimension with respect to \mathbb{R}^d serves as an extension of the Lebesgue dimension to discrete subsets by assigning a value between 0 and the Lebesgue dimension of the Euclidean space or infinity to each such subset. It further indicates that the lower Beurling dimension provides a subdivision for those sets which have infinite upper Beurling dimension. Moreover, for subsets with Beurling dimension, i.e., for which the lower Beurling dimension coincides with the upper Beurling dimension, it follows that this dimension is either 0, d , or ∞ . There are, however, examples of Λ , S and V not satisfying the conditions of Theorem 2.10 such that the Beurling dimension with respect to S is any non-negative number, as can be seen by considering the sequence $\Lambda = \{\pm n^\alpha : n \in \mathbb{N}\}$ with $S = \{0\}$ and $V = \mathbb{R}$.

We close this section by showing that Beurling dimension inherits the ‘‘intuitive’’ dimension of a discrete set.

Theorem 2.11. *Let $\Lambda \subset \mathbb{R}^d$ with $\mathcal{D}_d^+(\Lambda) < \infty$, and let $0 \leq n < d$. Suppose that Λ is contained in a translated bounded neighborhood of an n -dimensional subspace V of \mathbb{R}^d , i.e., $\Lambda \subset y_0 + \{x \in \mathbb{R}^d : \text{dist}(x, V) \leq \epsilon\}$ for $y_0 \in \mathbb{R}^d$ and $\epsilon > 0$. Then, for any affine subspace S such that there exists $z_0 \in \mathbb{R}^d$ satisfying $S + z_0 \supset V$, we have*

$$\dim_S^+(\Lambda) \leq n.$$

Proof. Let P be the orthogonal projection onto V . Note that $P(\Lambda)$ is an ϵ -perturbation of Λ , so by Theorem 2.7, $\dim_S^+(P(\Lambda)) = \dim_S^+(\Lambda)$. Moreover, since $S + z_0 \supset V$, Theorem 2.10 and Proposition 2.3 imply that $\dim_S^+(P(\Lambda)) \in [0, n] \cup \{\infty\}$, since $P(\Lambda)$ is contained in an n -dimensional subspace. Therefore, since $\dim_S^+(\Lambda) < \infty$, $\dim_S^+(\Lambda) = \dim_S^+(P(\Lambda)) \leq n$. \square

2.3. Comparison with other dimensions. In this section, we compare Beurling dimensions with other dimensions; namely, *mass dimensions*, \dim_{LM} and \dim_{UM} , and *discrete Hausdorff dimension*, \dim_H . The first can be defined as

$$\dim_{LM}(\Lambda) = \liminf_{n \rightarrow \infty} \frac{\ln \#(\Lambda \cap Q_n)}{\ln n}$$

and

$$\dim_{UM}(\Lambda) = \limsup_{n \rightarrow \infty} \frac{\ln \#(\Lambda \cap Q_n)}{\ln n},$$

see, e.g., [1]. (We shall see in this section that mass dimensions are special cases of Beurling dimensions.) For the precise definition of the *discrete Hausdorff dimension* we refer to [1]. For other notions of discrete dimensions and for a discussion of relations between them, we refer to [18] and [19].

First, we present yet another version of the definition of the Beurling dimension, which will facilitate comparisons to the mass dimensions.

Proposition 2.12. *Let $\Lambda \subset \mathbb{R}^d$ and S be an affine subspace of \mathbb{R}^d . Then we have*

$$\dim_S^-(\Lambda) = \liminf_{h \rightarrow \infty} \inf_{x \in S} \frac{\ln \#(\Lambda \cap Q_h(x))}{\ln h}$$

and

$$\dim_S^+(\Lambda) = \limsup_{h \rightarrow \infty} \sup_{x \in S} \frac{\ln \#(\Lambda \cap Q_h(x))}{\ln h}.$$

Proof. We only study the upper dimension. The proof for the lower dimension is similar.

First, consider $0 < r < \dim_S^+(\Lambda)$. In this case we have $\mathcal{D}_{S,r}^+(\Lambda) = \infty$. Therefore there exists a sequence $(h_n)_{n \in \mathbb{N}}$ with $\lim_{n \rightarrow \infty} h_n = \infty$ such that

$$\lim_{n \rightarrow \infty} \frac{\sup_{x \in S} \#(\Lambda \cap Q_{h_n}(x))}{h_n^r} = \infty.$$

Without loss of generality we can assume that for all $n \in \mathbb{N}$,

$$\frac{\sup_{x \in S} \#(\Lambda \cap Q_{h_n}(x))}{h_n^r} > 1.$$

This yields

$$\begin{aligned} \limsup_{h \rightarrow \infty} \sup_{x \in S} \frac{\ln \#(\Lambda \cap Q_h(x))}{\ln h} &\geq \limsup_{n \rightarrow \infty} \sup_{x \in S} \frac{\ln \#(\Lambda \cap Q_{h_n}(x))}{\ln h_n} \\ &> \limsup_{n \rightarrow \infty} \frac{\ln(h_n^r)}{\ln h_n} = r. \end{aligned}$$

Since this holds for every $r < \dim_S^+(\Lambda)$, we obtain

$$(3) \quad \limsup_{h \rightarrow \infty} \sup_{x \in S} \frac{\ln \#(\Lambda \cap Q_h(x))}{\ln h} \geq \dim_S^+(\Lambda).$$

Secondly, let $r > \dim_S^+(\Lambda)$, which implies $\mathcal{D}_{S,r}^+(\Lambda) = 0$. Fix $\epsilon > 0$. Then there exists $H > 0$ such that for all $h > H$:

$$\frac{\sup_{x \in S} \#(\Lambda \cap Q_h(x))}{h^r} \leq \epsilon.$$

Therefore we obtain

$$\limsup_{h \rightarrow \infty} \sup_{x \in S} \frac{\ln \#(\Lambda \cap Q_h(x))}{\ln h} \leq \limsup_{h \rightarrow \infty} \frac{\ln(\epsilon h^r)}{\ln h} = \limsup_{h \rightarrow \infty} \frac{\ln \epsilon + r \ln h}{\ln h} = r.$$

Thus

$$(4) \quad \limsup_{h \rightarrow \infty} \sup_{x \in S} \frac{\ln \#(\Lambda \cap Q_h(x))}{\ln h} \leq \dim_S^+(\Lambda).$$

Now (3) and (4) yield the claim. \square

Proposition 2.12 implies that mass dimension is Beurling dimension with respect to $S = \{0\}$. We state it formally in the following corollary, where \dim_{LM} and \dim_{UM} denote the lower and upper mass dimensions, respectively.

Corollary 2.13. *For every $\Lambda \subset \mathbb{R}^d$ and S an affine subspace of \mathbb{R}^d ,*

$$\dim_S^-(\Lambda) \leq \dim_{LM}(\Lambda) = \dim_{\{0\}}^-(\Lambda) \leq \dim_{UM}(\Lambda) = \dim_{\{0\}}^+(\Lambda) \leq \dim_S^+(\Lambda).$$

Proof. This follows immediately from Proposition 2.12 and the definition of mass dimension. \square

We present now several examples which illustrate the differences between Beurling dimension with respect to $S = \mathbb{R}^d$, mass dimensions, and (to a lesser extent) discrete Hausdorff dimension.

Example 2.14. *Let $\Lambda = \{(m, n) : m \in \mathbb{N}, n \in \mathbb{Z}\} \subset \mathbb{R}^2$. Then, $\dim_{\mathbb{R}^d}^-(\Lambda) = 0$ and $\dim_{\mathbb{R}^d}^+(\Lambda) = 2$, whereas we have $\dim_{LM}(\Lambda) = \dim_{UM}(\Lambda) = 2$.*

Example 2.15. *Define the set $\Lambda \subset \mathbb{Z}^2$ to be the union of sets $\Lambda_n, n \in \mathbb{N}$, where $\Lambda_n = \{(k, l) : 2^n \leq k < 2^n + 2^{n-1}, 0 \leq l < 2^{n-1}\}$. Then, it is not difficult to observe that $\dim_{\mathbb{R}^d}^-(\Lambda) = 0$ and $\dim_{\mathbb{R}^d}^+(\Lambda) = 2$, since the set Λ contains arbitrarily large pieces of the lattice \mathbb{Z}^2 . On the other hand, the mass dimension of Λ exists and is equal to 1, and the discrete Hausdorff dimension of Λ is equal to 0.*

Example 2.16. *Define the set $\Lambda \subset \mathbb{R}^2$ as follows: for each $n \in \mathbb{N}$ define $\Lambda_n = \{(2^n - k/2^n, l/2^n) : 1 \leq k, l \leq 2^n\}$; let $\Lambda = \bigcup_{n=1}^{\infty} \Lambda_n$. It follows immediately from the definition that the upper Beurling dimension of Λ with respect to $S = \mathbb{R}^d$ is infinity, because of the increasing concentration of points. On the other hand, it is not difficult to observe that the upper mass dimension of Λ satisfies $\dim_{UM}(\Lambda) = 2$. Moreover, the discrete Hausdorff dimension of any set is estimated from above by the upper mass dimension, and so $\dim_H(\Lambda) \leq 2$.*

3. APPLICATION TO PSEUDOFrames FOR SUBSPACES

We start by stating the definition of a pseudoframe for subspaces (see [16]). In the following, given a closed subspace E of a separable Hilbert space, we will always denote the orthogonal projection onto this subspace by P_E .

Definition 3.1. *Let E be a closed subspace of a separable Hilbert space \mathcal{H} , and let $\{x_i\}_{i \in I}$ be a sequence in \mathcal{H} . Then $\{x_i\}_{i \in I}$ is a Bessel sequence with respect to E , if*

$$\sum_{i \in I} |\langle f, x_i \rangle|^2 < \infty \quad \text{for all } f \in E.$$

A Bessel sequence $\{x_i\}_{i \in I}$ w.r.t. E is called a pseudoframe for the subspace E (PFFS for E), if there exists another Bessel sequence $\{x_i^*\}_{i \in I}$ in \mathcal{H} such that

$$f = \sum_{i \in I} \langle f, x_i \rangle x_i^* \quad \text{for all } f \in E.$$

The collection $\{x_i^*\}_{i \in I}$ is called a dual pseudoframe of $\{x_i\}_{i \in I}$ for the subspace E .

A Bessel sequence $\{x_i\}_{i \in I}$ w.r.t. E is called a pseudo-Riesz sequence for the subspace E (PRFS for E), if $\{P_E x_i\}_{i \in I}$ is a Riesz sequence in E .

In the sequel we will make use of the following characterization of PFFS's.

Theorem 3.2. [16, Theorem 4] *Let E be a closed subspace of a separable Hilbert space \mathcal{H} , let $\{x_i\}_{i \in I} \subset \mathcal{H}$ be a Bessel sequence w.r.t. E , and let $\{x_i^*\}_{i \in I}$ be a Bessel sequence in \mathcal{H} . Then the following conditions are equivalent.*

- (i) $\{x_i\}_{i \in I}$ is a PFFS for E with dual pseudoframe $\{x_i^*\}_{i \in I}$.
- (ii) The following conditions hold.
 - (a) $\{P_E x_i\}_{i \in I}$ is a frame for E with dual frame $\{P_E x_i^*\}_{i \in I}$.
 - (b) For all $f \in H$, $\sum_{i \in I} \langle f, P_E x_i \rangle (I - P_E) x_i^* = 0$.

Note that in the case that $\{P_E x_i\}_{i \in I}$ is a frame for the subspace E , one can always find $\{x_i^*\}_{i \in I} \subset E$ satisfying the conditions for PFFS. Thus, for our purposes, we will not refer to the dual frame, and we write briefly that $\{x_i\}_{i \in I}$ is a PFFS for E . Furthermore, we say that the *frame bounds of a PFFS for E* are the frame bounds of the frame $\{P_E x_i : i \in I\}$ for E .

In this section, we consider PFFS's with more structure than the general case. In the following let $\mathcal{G} = \{g_1, \dots, g_K\}$ be some finite collection of functions in $L^2(\mathbb{R}^d)$, let $\Lambda = \{\Lambda_1, \dots, \Lambda_K\}$ be a finite collection of subsets of \mathbb{R}^{2d} , and let the associated *Gabor system* be defined by

$$\begin{aligned} \mathcal{G}(G, \Lambda) &= \{e^{2\pi i x \cdot t} g_k(t - y) : (x, y) \in \Lambda_k, 1 \leq k \leq K\} \\ &= \{M_x T_y g_k : (x, y) \in \Lambda_k, 1 \leq k \leq K\}, \end{aligned}$$

where M_x and T_y are the modulation and translation operators respectively. The discrete set Λ will be referred to as the *set of parameters*. We will consider those Gabor systems which are PFFS's for $L^2(E)$ for some $E \subset \mathbb{R}^d$. When E is, for example, a bounded set, one would expect that the modulations and translations together would need to be "sufficiently dense" in order to form a PFFS for $L^2(E)$. One would also

expect that the collection of modulations and translations needs to be “sufficiently sparse” in order to form a PRFS. We make these intuitive notions precise in Theorem 3.3 below.

The sets E we will consider will be of the following form. For $E \subset \mathbb{R}^d$ and $0 \leq m \leq d$, we say E contains a tube around an m -dimensional space if there exists an affine subspace A of \mathbb{R}^d of dimension m and an $\epsilon > 0$ such that $\{z \in \mathbb{R}^d : \text{dist}(z, A) < \epsilon\} \subset E$.

With this notation, we are ready to state our main theorem, whose proof will be given at the end of this section.

Theorem 3.3. *Let $\mathcal{G} = \{g_1, \dots, g_K\} \subset L^2(\mathbb{R}^d) \setminus \{0\}$, and let $\Lambda = \{\Lambda_1, \dots, \Lambda_K\}$ be a finite collection of subsets of \mathbb{R}^{2d} . Furthermore, let E be a subset of \mathbb{R}^d , which contains a tube around an m -dimensional affine subspace A of \mathbb{R}^d .*

- (i) *If $\mathcal{G}(G, \Lambda)$ is a PFFS for $L^2(E)$, then either $\dim_{\mathbb{R}^d \times A}^-(\Lambda) \geq d + m$ or Λ is not relatively uniformly separated.*
- (ii) *If $\mathcal{G}(G, \Lambda)$ is a PRFS for $L^2(E)$, then $\dim_{\mathbb{R}^d \times A}^-(\Lambda) = \dim_{\mathbb{R}^d \times A}^+(\Lambda) = d + m$.*

Let us briefly compare this result with Nyquist density results of Gabor systems. For Gabor systems, the following result from [5] holds.

Theorem 3.4. [5, Theorem 1.1 and 3.6] *Let $\mathcal{G} = \{g_1, \dots, g_K\} \subset L^2(\mathbb{R}^d) \setminus \{0\}$, and let $\Lambda \subset \mathbb{R}^{2d}$.*

- (i) *If $\mathcal{G}(G, \Lambda)$ is a frame for $L^2(\mathbb{R}^d)$, then $\mathcal{D}_{\mathbb{R}^{2d}, d}^-(\Lambda) \geq 1$.*
- (ii) *If $\mathcal{G}(G, \Lambda)$ is a Riesz basis for $L^2(\mathbb{R}^d)$, then $\mathcal{D}_{\mathbb{R}^{2d}, d}^-(\Lambda) = \mathcal{D}_{\mathbb{R}^{2d}, d}^+(\Lambda) = 1$.*

It can be easily seen that in the situation of Gabor PFFS’s Beurling dimension plays the role which Beurling density plays for Gabor frames. It is in this sense that it is justified to regard the Beurling dimension as a *Nyquist dimension*.

Note also that, in particular, Theorem 3.3 implies that if a collection of modulations is a frame for $L^2([0, 1])$, then the modulations must have positive upper Beurling density, recovering a result of Christensen, Deng, and Heil [5].

In general, for a PFFS $\mathcal{G}(G, \Lambda)$, if Λ is not relatively separated (in particular, $\mathcal{G}(G, \Lambda)$ is not Bessel in $L^2(\mathbb{R}^d)$), then the Λ can be quite odd. For example, in the case that $E \subset \mathbb{R}$, there is no restriction on those $(x, y) \in \Lambda$ for which the support of $T_x g$ is disjoint from E for PFFS’s, so the lower dimension can be made as large as desired. It is also possible to construct examples of PFFS’s for $L^2(E)$ for which $\dim_{\mathbb{R}^d \times A}^-(\Lambda)$ is less than $d + m$, see Example 3.13. It is thus more surprising that in the case of PRFS’s, we are able to obtain a Nyquist dimension as in Theorem 3.3 (ii).

Our method of proof is inspired by techniques to prove density results given in the recent preprint [10]. We will show that PFFS’s for $L^2(E)$ of the type mentioned in Theorem 3.3 satisfy a modified version of the Homogeneous Approximation Property (HAP) of Ramanathan and Steger. Using this modified HAP, we will then show that the set of parameters Λ has lower dimension greater than or equal to $d + m$. We will also use the results from Section 2 to show that the upper dimension of a PRFS is bounded above by $d + m$.

3.1. Preliminary Lemmas. We begin by recalling the following definition and lemma from [10].

Definition 3.5. (i) Given a set $U \subset \mathbb{R}^{2d}$, for each $t \geq 0$ define

$$U_t = \{x \in \mathbb{R}^{2d} : \text{dist}(x, U) < t\}.$$

(ii) The Fréchet distance between two closed sets $U, V \subset \mathbb{R}^{2d}$ is

$$[U, V] = \inf\{t \geq 0 : U \subset V_t \text{ and } V \subset U_t\}.$$

(iii) Given closed sets $U_n \subset \mathbb{R}^{2d}$ and given a closed set $V \subset \mathbb{R}^{2d}$, we say that U_n converges weakly to V if

$$\lim_{n \rightarrow \infty} [U_n \cap K, V \cap K] = 0 \quad \text{for all compact } K \subset \mathbb{R}^{2d}.$$

In this case, we write $U_n \rightarrow^w V$.

Lemma 3.6. [10, Lemma 2.10] Let $\Lambda \subset \mathbb{R}^{2d}$ be a countable sequence which is δ -uniformly separated for some $\delta > 0$. Then given any sequence $\{z_n\}_{n \in \mathbb{N}}$ of points in \mathbb{R}^{2d} , there exists a subsequence $\{w_n\}_{n \in \mathbb{N}}$ of $\{z_n\}_{n \in \mathbb{N}}$ and a sequence $\Lambda' \subset \mathbb{R}^{2d}$ such that

$$\Lambda - w_n \rightarrow^w \Lambda' \text{ as } j \rightarrow \infty, \quad k = 1, \dots, N.$$

The following two lemmas will be heavily employed in the proof of Theorem 3.10.

Lemma 3.7. Let $a \leq d$ be a nonnegative integer, $G = \{g_1, \dots, g_K\} \subset L^2(\mathbb{R}^d) \setminus \{0\}$, $\Lambda = \{\Lambda_1, \dots, \Lambda_K\}$ be a finite collection of subsets of \mathbb{R}^{2d} , and $E \subset \mathbb{R}^d$ be such that the following conditions hold.

- (a) Each Λ_k is 2δ -uniformly separated for some $\delta > 0$.
- (b) E is a tube around the subspace $\{x \in \mathbb{R}^d : x_{d-a+1} = \dots = x_d = 0\}$ for $1 \leq a \leq d$, or E is a convex neighborhood of the origin if $a = 0$.
- (c) $\mathcal{G}(G, \Lambda)$ is a PFFS for $L^2(E)$ with frame bounds A and B .

Then, for each $x_0 \in \mathbb{R}^d$, $y_0 \in \mathbb{R}^a$, and $0 \in \mathbb{R}^{d-a}$, $\{M_x T_y g_k : (x, y) \in \Lambda_k + (x_0, y_0, 0) : 1 \leq k \leq K\}$ is a PFFS for $L^2(E)$ with frame bounds A and B .

Proof. Let $h \in L^2(E)$, and define $\tilde{h}(t) = e^{-2\pi i x_0 t} h(t + (y_0, 0))$. Note that $\|h\|_2 = \|\tilde{h}\|_2$, $P_{L^2(E)} h = h$ and $P_{L^2(E)} \tilde{h} = \tilde{h}$. The result follows from the following computation:

$$\begin{aligned} \sum_{k=1}^K \sum_{(x,y) \in \Lambda_k + (x_0, y_0, 0)} |\langle P_{L^2(E)} M_x T_y g_k, h \rangle|^2 &= \sum_{k=1}^K \sum_{(x,y) \in \Lambda_k + (x_0, y_0, 0)} |\langle M_x T_y g_k, h \rangle|^2 \\ &= \sum_{k=1}^K \sum_{(x,y) \in \Lambda_k} |\langle M_x T_y g_k, \tilde{h} \rangle|^2 \\ &= \sum_{k=1}^K \sum_{(x,y) \in \Lambda_k} |\langle P_{L^2(E)} M_x T_y g_k, \tilde{h} \rangle|^2. \end{aligned}$$

□

Lemma 3.8. *Let a, E, G and $\Lambda = \{\Lambda_1, \dots, \Lambda_K\}$ be as in Lemma 3.7. Let $\{x_n\}_{n \in \mathbb{N}}$ be a sequence in \mathbb{R}^d , $\{y_n\}_{n \in \mathbb{N}}$ a sequence in \mathbb{R}^a and for each n let $z_n = (x_n, y_n, 0) \in \mathbb{R}^{2d}$. Suppose that for each $1 \leq k \leq K$, $\Lambda_k - z_n \xrightarrow{w} \Lambda'_k$. Then, $\mathcal{G}(G, \Lambda')$ is a PFFS for $L^2(E)$ with the same frame bounds, where $\Lambda' = \{\Lambda'_1, \dots, \Lambda'_K\}$.*

Proof. Note that for each $1 \leq k \leq K$, a cube of the form $Q_\delta(z)$ can contain at most one point in Λ_k . Using the weak convergence of $\Lambda_k - z_n \xrightarrow{w} \Lambda'_k$, it can also be shown that each cube $Q_\delta(z)$ can contain at most one point of Λ'_k .

Choose $\epsilon > 0$ and let $f \in M^1(\mathbb{R}^d) \cap L^2(E)$. Then, for each $1 \leq k \leq K$, the short-time Fourier transform $V_{g_k} f$ defined by $V_{g_k} f(x, y) = \langle f, M_x T_y g_k \rangle$, $(x, y) \in \mathbb{R}^d \times \mathbb{R}^d$ is an element of the amalgam space $W(L^\infty, \ell^2)$. By using an equivalent norm of this space, we can find an $m \in \mathbb{N}$ such that for each $1 \leq k \leq K$,

$$\sum_{j \in \mathbb{Z}^{2d} \setminus Q_m(0)} \|V_{g_k} f \cdot 1_{Q_\delta(\delta j)}\|_\infty^2 < \frac{\epsilon}{2}.$$

Fix $1 \leq k \leq K$. Set $R = (2m + 1)\delta$. If $(x, y) \in \mathbb{R}^{2d} \setminus Q_R(0)$, then there is a unique $j \in \mathbb{Z}^{2d} \setminus Q_m(0)$ such that $(x, y) \in Q_\delta(\delta j)$. Hence,

$$\begin{aligned} \sum_{(x,y) \in \Lambda'_k \setminus Q_R(0)} |\langle f, M_x T_y g_k \rangle|^2 &= \sum_{(x,y) \in \Lambda'_k \setminus Q_R(0)} |V_{g_k} f(x, y)|^2 \\ &\leq \sum_{j \in \mathbb{Z}^{2d} \setminus Q_m(0)} \sup_{(u,v) \in Q_\delta(\delta j)} |V_{g_k} f(u, v)|^2 < \frac{\epsilon}{2}. \end{aligned}$$

Similarly, for each $n \in \mathbb{N}$, we have

$$\sum_{(x,y) \in (\Lambda_k - z_n) \setminus Q_R(0)} |\langle f, M_x T_y g_k \rangle|^2 < \frac{\epsilon}{2}.$$

Let $D = \sup_{n \in \mathbb{N}} \#((\Lambda_k - z_n) \cap Q_R(0))$. We estimate the difference in the ℓ_2 norm of the inner products of f with $M_x T_y g_k$ in the cases that $(x, y) \in \Lambda'_k \cap Q_R(0)$ and $(x, y) \in \Lambda_k - z_n \cap Q_R(0)$. Clearly, when $D = 0$, the difference is 0, so we consider the case $D > 0$. Using the continuity of the modulation and translation operators and the fact that $Q_R(0)$ is compact, find $\theta > 0$ such that for all $(x, y) \in Q_R(0)$ and all $|u|, |v| < \theta$, $\|M_u T_v M_x T_y f - M_x T_y f\|_2 < \epsilon(2(ND)^{1/2} \|f\|_2)^{-1}$. Let n be large enough so that

$$[(\Lambda_k - z_n) \cap Q_R(0), \Lambda'_k \cap Q_R(0)] < \theta.$$

Then, for each $(x, y) \in (\Lambda_k - z_n) \cap Q_R(0)$, there exist unique points $|u(x, y)|, |v(x, y)| < \theta$ such that $(x + u(x, y), y + v(x, y)) \in \Lambda'_k$. Moreover, this correspondence yields a bijection between $(\Lambda_k - z_n) \cap Q_R(0)$ and $\Lambda_k \cap Q_R(0)$. For notational ease, we write $\Lambda_{n,R} = (\Lambda_k - z_n) \cap Q_R(0)$, where the dependence on k is suppressed since k is fixed.

We compute

$$\begin{aligned}
& \left| \left(\sum_{(x,y) \in \Lambda'_k \cap Q_R(0)} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} - \left(\sum_{(x,y) \in \Lambda_{n,R}} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} \right| \\
& \leq \left(\sum_{(x,y) \in \Lambda_{n,R}} |\langle f, M_{x+u(x,y)} T_{y+v(x,y)} g_k - M_x T_y g_k \rangle|^2 \right)^{1/2} \\
& \leq \left(\sum_{(x,y) \in \Lambda_{n,R}} \|f\|^2 \|M_{x+u(x,y)} T_{y+v(x,y)} g_k - M_x T_y g_k\|_2^2 \right)^{1/2} < \frac{\epsilon}{2}.
\end{aligned}$$

Now let A and B denote the frame bounds of $\mathcal{G}(G, \Lambda)$, and Recall that $z_n = (x_n, y_n, 0)$, where $y_n \in \mathbb{R}^a$ and $0 \in \mathbb{R}^{d-a}$. Employing (b), for each $f \in L^2(E)$ we have $M_{x_n} T_{(y_n, 0)} f \in L^2(E)$. Therefore, $\langle M_{x_n} T_{(y_n, 0)} f, P_{L^2(E)} g_k \rangle = \langle M_{x_n} T_{(y_n, 0)} f, g_k \rangle$. In particular, $\|P_{L^2(E)} M_{x_n} T_{(y_n, 0)} f\|_2 = \|f\|_2$. With this observation, we allow k to vary again and we compute

$$\begin{aligned}
& \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda'_k} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} \\
& \leq \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda'_k \cap Q_R(0)} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} + \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda'_k \setminus Q_R(0)} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} \\
& \leq \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda'_k \cap Q_R(0)} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} + \frac{\epsilon}{2} + \frac{\epsilon}{2} \\
& \leq \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda_k} |\langle M_{x_n} T_{(y_n, 0)} f, M_x T_y g_k \rangle|^2 \right)^{1/2} + \epsilon \\
& \leq B^{1/2} \|M_{x_n} T_{(y_n, 0)} f\|_2 + \epsilon \\
& = B^{1/2} \|f\|_2 + \epsilon.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
& \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda'_k} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} \\
& \geq \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda'_k \cap Q_R(0)} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} \\
& \geq \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda'_k \cap Q_R(0)} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2}
\end{aligned}$$

$$\begin{aligned}
&\geq \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda_k - z_n \cap Q_R(0)} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} - \frac{\epsilon}{2} \\
&\geq \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda_k - z_n} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} - \left(\sum_{k=1}^K \sum_{(x,y) \in \Lambda_k - z_n \setminus Q_R(0)} |\langle f, M_x T_y g_k \rangle|^2 \right)^{1/2} - \frac{\epsilon}{2} \\
&\geq A^{1/2} \|f\|_2 - \frac{\epsilon}{2} - \frac{\epsilon}{2} \\
&= A^{1/2} \|f\|_2 - \epsilon.
\end{aligned}$$

In the last line, we have used Lemma 3.7. Since $\epsilon > 0$ is arbitrary, $\{P_{L^2(E)} M_x T_y g_k : (x, y) \in \Lambda'_k, 1 \leq k \leq K\}$ is a frame for $L^2(E)$, i.e. $\mathcal{G}(G, \Lambda')$ is a PFFS for $L^2(E)$ with frame bounds A and B . \square

It is clear that Lemma 3.8 is not true for general z_n . Indeed, consider $E = [0, 1]$, $g = 1_E$, $\Lambda = \mathbb{Z} \times \{0\}$. Then $\{M_x T_y g : (x, y) \in \Lambda\}$ is a PFFS for $L^2(E)$, but $\Lambda + (0, 1) = \mathbb{Z} \times \{1\}$ is not.

3.2. The A-Ramanathan-Steger Weak Homogeneous Approximation Property. The Homogeneous Approximation Property (HAP) (cf. [5]) is a common tool to study density conditions of Gabor systems. However, it is not difficult to see that PFFS's do not generally satisfy the HAP. Therefore, we now define a weaker notion of the HAP which PFFS's do satisfy, thereby deriving interesting approximation properties of PFFS.

Definition 3.9. Let $E \subset \mathbb{R}^d$. For $h > 0$ and $(u, \eta) \in \mathbb{R}^{2d}$, we set $W(h, u, \eta) = \text{span}\{M_x T_y P_{L^2(E)} g : (x, y) \in \Lambda \cap Q_h(u, \eta)\}$. Let $A \subset \mathbb{R}^{2d}$. We say that the PFFS for $L^2(E)$ $\mathcal{G}(G, \Lambda)$ possesses the Ramanathan-Steger Weak Homogeneous Approximation Property with respect to A if for all $f \in L^2(E)$ and for all $\epsilon > 0$, there exists $R > 0$ such that for all $(u, \eta) \in A$, $\text{dist}(M_\eta T_u f, W(R, u, \eta)) < \epsilon$.

For this paper, we will abbreviate the Ramanathan-Steger Weak Homogeneous Approximation Property with respect to A by simply the *weak HAP with respect to A* .

Theorem 3.10. Let $a \leq d$ be a nonnegative integer, $G = \{g_1, \dots, g_K\} \subset L^2(\mathbb{R}^d) \setminus \{0\}$, $\Lambda \subset \mathbb{R}^{2d}$ and $E \subset \mathbb{R}^d$ be such that

- (a) There exists $\delta > 0$ such that for $1 \leq k \leq K$, Λ_k is 2δ -uniformly separated,
- (b) E is a tube around the subspace $\{x \in \mathbb{R}^d : x_{d-a+1} = \dots = x_d = 0\}$ for $1 \leq a \leq d$, or E is a convex neighborhood of the origin if $a = 0$, and
- (c) $\mathcal{G}(G, \Lambda)$ is a PFFS for $L^2(E)$.

Let $A \subset \mathbb{R}^d \times \mathbb{R}^{d-a} \times \{0\}^a$. Then, $\mathcal{G}(G, \Lambda)$ possesses the weak HAP with respect to A .

Proof. Suppose that the weak HAP with respect to A fails. Then, there exists a function $f \in L^2(E)$ and $\epsilon > 0$ such that for each $n \in \mathbb{N}$, there exists $z_n = (u_n, \eta_n) \in A$

such that $\text{dist}(M_{\eta_n}T_{u_n}f, W(n, u_n, \eta_n)) > \epsilon$. By Lemma 3.6, there exists a subsequence $\{w_n\}$ of $\{z_n\}$ and Λ'_k such that for each $1 \leq k \leq K$, we have

$$\Lambda_k - w_n \xrightarrow{w} \Lambda'_k \text{ as } n \rightarrow \infty.$$

Therefore, by Lemma 3.8, $\mathcal{G}(G, \Lambda')$ is a PFFS for $L^2(E)$ with the same frame bounds. We claim now that for any $R > 0$,

$$(5) \quad \text{dist}\left(f, \text{span}\{M_\xi T_x P_{L^2(E)} g_k : (x, \xi) \in \Lambda'_k \cap Q_R(0), k = 1, \dots, K\}\right) \geq \frac{\epsilon}{2}.$$

To see this, choose any scalars $\{c_{k,x,\xi} : (x, \xi) \in \Lambda'_k \cap Q_R(0), k = 1, \dots, K\}$. Let

$$D = \sum_{k=1}^K \sum_{(x,\xi) \in \Lambda'_k \cap Q_R(0)} |c_{k,x,\xi}|.$$

From above, we know that $\|f - 0\|_2 \geq \epsilon$, so we may assume that $D \neq 0$. Since the family of modulation-translation operators is strongly continuous (cf. [10, Lemma 2.1]), there exists $\theta < \delta/2$ such that whenever $|x|, |\xi| < \theta$, $(u, \eta) \in A$ and $1 \leq k \leq K$,

$$\|M_{\xi+\eta}T_{x+u}P_{L^2(E)}g_k - M_\xi T_x P_{L^2(E)}g_k\|_2 < \frac{\epsilon}{2D}.$$

As in the proof of Lemma 3.8, we can find n large enough so that each point of $\Lambda'_k \cap Q_R(0)$ is within θ of a unique point in $\Lambda_k - w_n \cap Q_R(0)$, and conversely. So, we can write

$$\Lambda_k - w_n \cap Q_R(0) = \{(x + u(x, \xi, k), \xi + \eta(x, \xi, k)) : (x, \xi) \in \Lambda'_k \cap Q_R(0), k = 1, \dots, K\},$$

with $|u(x, \xi, k)|, |\eta(x, \xi, k)| < \theta$. Hence,

$$\begin{aligned} & \left\| f - \sum_{k=1}^K \sum_{(x,\xi) \in \Lambda'_k \cap Q_R(0)} c_{k,x,\xi} M_\xi T_x P_{L^2(E)} g_k \right\|_2 \\ & \geq \left\| f - \sum_{k=1}^K \sum_{(x,\xi) \in \Lambda_k - w_n \cap Q_R(0)} c_{k,x,\xi} M_\xi T_x P_{L^2(E)} g_k \right\|_2 \\ & \quad - \left\| \sum_{k=1}^K \sum_{(x,\xi) \in \Lambda'_k \cap Q_R(0)} c_{k,x,\xi} (M_{\xi+\eta(x,\xi,k)} T_{x+u(x,\xi,k)} P_{L^2(E)} g_k - M_\xi T_x P_{L^2(E)} g_k) \right\|_2 \\ & \geq \text{dist}\left(f, \text{span}\{M_\xi T_x P_{L^2(E)} g_k : (x, \xi) \in \Lambda_k - w_n \cap Q_R(0), k = 1, \dots, K\}\right) \\ & \quad - \sum_{k=1}^K \sum_{(x,\xi) \in \Lambda'_k \cap Q_R(0)} |c_{k,x,\xi}| \|M_{\xi+\eta(x,\xi,k)} T_{x+u(x,\xi,k)} P_{L^2(E)} g_k - M_\xi T_x P_{L^2(E)} g_k\|_2 \\ & \geq \epsilon - \sum_{k=1}^K \sum_{(x,\xi) \in \Lambda'_k \cap Q_R(0)} |c_{k,x,\xi}| \frac{\epsilon}{2D} = \frac{\epsilon}{2}. \end{aligned}$$

Since this is true for every choice of scalars, we conclude that (5) holds. But since R is arbitrary, this implies that $f \notin \overline{\text{span}}(P_{L^2(E)}\mathcal{G}(G, \Lambda'))$. Therefore, there exists some $0 \neq h \in L^2(E) \cap (\overline{\text{span}}(P_{L^2(E)}\mathcal{G}(G, \Lambda')))^{\perp}$. This contradicts the fact that $P_{L^2(E)}\mathcal{G}(G, \Lambda')$ is a frame for $L^2(E)$ (hence complete in $L^2(E)$). \square

Employing the weak HAP as a main ingredient for the proof, we will show that sets of parameters of PFFS's are comparable with sets of parameters of Riesz sequences by means of their densities $D_{A,r}^{\pm}$ (recall Definition 2.1).

Theorem 3.11. *Let a, E, G, A and Λ be as in Theorem 3.10. Let $\Delta_1, \dots, \Delta_L \subset A$ and $\phi_1, \dots, \phi_L \in L^2(E) \setminus \{0\}$. Assume that $\Phi = P_{L^2(E)}\mathcal{G}(\phi_1, \dots, \phi_L, \Delta_1, \dots, \Delta_L)$ is a Riesz sequence in $L^2(E)$. Let $\Lambda = \cup_{k=1}^K \Lambda_k$ and $\Delta = \cup_{k=1}^L \Delta_k$. Then $D_{A,r}^+(\Lambda) \geq D_{A,r}^+(\Delta)$ and $D_{A,r}^-(\Lambda) \geq D_{A,r}^-(\Delta)$ for all $0 < r < \infty$.*

Proof. Note that by Theorem 3.10 we have that $\mathcal{G} = \mathcal{G}(G, \Lambda)$ possesses the weak HAP with respect to A . Let

$$\tilde{\mathcal{G}} = \bigcup_{k=1}^K \{\tilde{g}_{x,\xi,k}\}_{(x,\xi) \in \Lambda_k}$$

denote the canonical dual frame of $P_{L^2(E)}\mathcal{G}$ in $L^2(E)$. Let

$$\tilde{\Phi} = \bigcup_{k=1}^L \{\tilde{\phi}_{x,\xi,k}\}_{(x,\xi) \in \Delta_k}$$

denote the dual frame within the closed linear span of Φ .

Given $h > 0$ and $(u, \eta) \in A$, set

$$\begin{aligned} \tilde{W}(h, u, \eta) &= \text{span}\{M_{\xi}T_x\tilde{g}_k : (x, \xi) \in \Lambda_k \cap Q_h(u, \eta), k = 1, \dots, K\} \\ V(h, u, \eta) &= \text{span}\{M_{\xi}T_x\phi_k : (x, \xi) \in \Delta_k \cap Q_h(u, \eta), k = 1, \dots, L\}. \end{aligned}$$

Since we have assumed that each Λ_k is uniformly separated, we have that $D_{\mathbb{R}^d, r}^+(\Lambda_k) < \infty$. So, \tilde{W} is a finite dimensional space.

Fix $\epsilon > 0$. Applying the definition of weak HAP with respect to A to the functions $f = \phi_k$, we see that there exists an $R > 0$ such that

$$(6) \quad \text{dist}(M_{\eta}T_u\phi_k, \tilde{W}(R, u, \eta)) < \frac{\epsilon}{D}, \quad \text{for all } (u, \eta) \in A, k = 1, \dots, L,$$

where

$$D = \sup\{\|\tilde{\phi}_{u,\eta,k}\| : (u, \eta) \in \Delta_k, k = 1, \dots, L\}.$$

Fix an $h > 0$ and $(u, \eta) \in \mathbb{R}^{2d}$. For simplicity, set $V = V(h, u, \eta)$ and $W = \tilde{W}(R + h, u, \eta)$. Define $T : V \rightarrow V$ by $T = P_V P_W$. Note that T is self-adjoint and W is finite-dimensional, so T has a finite, real trace.

We now estimate the trace of T . An easy upper bound is given by

$$(7) \quad \text{trace}(T) \leq \text{rank}(T) \leq \dim(W) = \#(\Lambda \cap Q_{R+h}(u, \eta)).$$

For a lower bound, note that $\{P_{L^2(E)}M_\xi T_x \phi_k : (x, \xi) \in \Delta_k \cap Q_h(u, \eta), k = 1, \dots, L\}$ is a basis for the finite-dimensional space V . The dual basis in V is the biorthogonal system in V , which is

$$\{P_V \tilde{\phi}_{x, \xi, k} : (x, \xi) \in \Delta_k \cap Q_h(u, \eta), k = 1, \dots, L\}.$$

Therefore, using that $TP_{L^2(E)} = T$, we compute

$$\begin{aligned} \text{trace}(T) &= \sum_{k=1}^L \sum_{(x, \xi) \in \Delta_k \cap Q_h(u, \eta)} \langle T(P_{L^2(E)}M_\xi T_x \phi_k), P_V \tilde{\phi}_{x, \xi, k} \rangle \\ &= \sum_{k=1}^L \sum_{(x, \xi) \in \Delta_k \cap Q_h(u, \eta)} \left(\langle M_\xi T_x \phi_k, P_V \tilde{\phi}_{x, \xi, k} \rangle + \langle (P_W - I)(M_\xi T_x \phi_k), P_V \tilde{\phi}_{x, \xi, k} \rangle \right) \\ &= \sum_{k=1}^L \sum_{(x, \xi) \in \Delta_k \cap Q_h(u, \eta)} (1 - \langle (P_W - I)(M_\xi T_x \phi_k), P_V \tilde{\phi}_{x, \xi, k} \rangle). \end{aligned}$$

Additionally, if $(x, \xi) \in \Delta \cap Q_h(u, \eta)$, then we have $Q_R(x, \xi) \subset Q_{R+h}(u, \eta)$, hence $W(R, x, \xi) \subset W(R+h, u, \eta)$ and therefore

$$\begin{aligned} |\langle (P_W - I)(M_\xi T_x \phi_k), P_V \tilde{\phi}_{x, \xi, k} \rangle| &\leq \|(P_W - I)(M_\xi T_x \phi_k)\|_2 \|P_V \tilde{\phi}_{x, \xi, k}\| \\ &\leq \text{dist}(M_\xi T_x \phi_k, W(R, x, \xi)) \|\tilde{\phi}_{x, \xi, k}\|_2 \\ &\leq \frac{\epsilon}{D} D = \epsilon. \end{aligned}$$

Therefore, we have that

$$\text{trace}(T) \geq \sum_{k=1}^L \sum_{(x, \xi) \in \Delta_k \cap Q_h(u, \eta)} (1 - \epsilon) = (1 - \epsilon) \#(\Delta \cap Q_h(u, \eta)).$$

Finally, combining our upper and lower estimates yields

$$(1 - \epsilon) \#(\Delta \cap Q_h(u, \eta)) \leq \#(\Lambda \cap Q_{R+h}(u, \eta)), \quad \text{for all } (u, \eta) \in A, h > 0,$$

and so for each $r > 0$,

$$\begin{aligned} D_{A,r}^+(\Delta) &= \limsup_{h \rightarrow \infty} \inf_{(u, \eta) \in A} \frac{\#(\Delta \cap Q_h(u, \eta))}{h^r} \\ &\leq \frac{1}{1 - \epsilon} \limsup_{h \rightarrow \infty} \sup_{(u, \eta) \in A} \frac{\#(\Lambda \cap Q_{R+h}(u, \eta)) (R+h)^r}{(R+h)^r h^r} \\ &= \frac{1}{1 - \epsilon} D_{A,r}^+(\Lambda). \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we conclude that $D_{A,r}^+(\Delta) \leq D_{A,r}^+(\Lambda)$, and a similar calculation shows that $D_{A,r}^-(\Delta) \leq D_{A,r}^-(\Lambda)$. \square

3.3. Beurling Dimension of Gabor PFFS's and PRFS's. The following result gives a detailed account of the upper and lower dimensions of sets of parameters of PFFS's and PRFS's for an arbitrary subset $A \subset \mathbb{R}^d \times \mathbb{R}^{d-a} \times \{0\}^a$. This will give rise to Theorem 3.3 by choosing an appropriate subset A .

Theorem 3.12. *Let $1 \leq a \leq d$ be an integer, $G = \{g_1, \dots, g_K\} \subset L^2(\mathbb{R}^d) \setminus \{0\}$, $\{\Lambda_1, \dots, \Lambda_K\}$ be a collection of subsets of \mathbb{R}^{2d} and $E \subset \mathbb{R}^d$ be such that*

- (a) *E is a tube around the subspace $\{x \in \mathbb{R}^d : x_{d-a+1} = \dots = x_d = 0\}$, and*
- (b) *$\mathcal{G}(G, \Lambda)$ is a PFFS for $L^2(E)$.*

Further, let $S = \mathbb{R}^d \times \mathbb{R}^{d-a} \times \{0\}^a$. Then the following conditions hold.

- (i) *Let F contain E . If $\mathcal{G}(G, \Lambda)$ is a PFFS for $L^2(F)$, then for $\Lambda = \cup_{k=1}^K \Lambda_k$,*

$$\dim_{\bar{S}}^-(\Lambda) \geq 2d - a \text{ or } \dim_{\mathbb{R}^d}^+(\Lambda) = \infty.$$

In particular, if $\mathcal{G}(G, \Lambda)$ is a Bessel sequence in $L^2(\mathbb{R}^d)$, then $\dim_{\bar{S}}^-(\Lambda) \geq 2d - a$.

- (ii) *Let H be contained in E . If $\mathcal{G}(G, \Lambda)$ is a PRFS for $L^2(H)$, then*

$$\dim_{\bar{S}}^-(\Lambda) = \dim_{\bar{S}}^+(\Lambda) = 2d - a.$$

Proof. (i): Suppose for each $1 \leq k \leq K$, $\dim_{\mathbb{R}^d}^+(\Lambda_k) < \infty$. Then, by Proposition 2.9 and repeating some Λ_k 's if necessary, we may assume that each Λ_k is uniformly separated. Note that if $E = \{(x_1, \dots, x_d) \in \mathbb{R}^d : |x_i| \leq \epsilon \text{ for } d - a + 1 \leq i \leq d \text{ and some } \epsilon > 0\}$, then let $\phi = 1_{[-\epsilon, \epsilon]}$ and $\Delta = \frac{1}{2\epsilon} \mathbb{Z}^d \times 2\epsilon \mathbb{Z}^{d-a} \times \{0\}^a$ and observe that $\{M_x T_y \phi : (x, y) \in \Delta\}$ is an ONB for $L^2(E)$. Moreover,

$$D_{S,r}^+(\Delta) = D_{S,r}^-(\Delta) = \begin{cases} \infty & : 0 \leq r < 2d - a, \\ \frac{(2\epsilon)^{d-a}}{(2\epsilon)^d} & : r = 2d - a, \\ 0 & : r > 2d - a. \end{cases}$$

Therefore, by Theorem 3.11, if $\mathcal{G}(G, \Lambda)$ is a PFFS for $L^2(E)$, then $\dim_{\bar{S}}^-(\Lambda) \geq 2d - a$.

Moreover, if F merely contains $E = \{(x_1, \dots, x_d) \in \mathbb{R}^d : |x_i| \leq \epsilon \text{ for } d - a + 1 \leq i \leq d \text{ and some } \epsilon > 0\}$, then since the projection of a frame is a frame, the result follows from the case considered in the first part of this proof.

(ii): Clearly, it suffices to prove (ii) in the case that $K = 1$. First, note that by applying Theorem 3.11 to compare the PRFS given in the statement of Theorem 3.12 to the frame for $L^2(\mathbb{R}^d)$ given by $\mathcal{G}(1_{[0,1]^d}, \mathbb{Z}^{2d})$, it follows that for all $0 < r < \infty$, $D_{\mathbb{R}^d, r}^+(\Lambda) \leq D_{\mathbb{R}^d, r}^-(\mathbb{Z}^{2d})$. In particular, by part (i), this implies that $\dim_{\mathbb{R}^{2d}}(\Lambda) < \infty$, so $\dim_{\bar{S}}^-(\Lambda) \geq 2d - a$.

Now, we will show that $\dim_{\bar{S}}^+(\Lambda) \leq 2d - a$. We will actually show the stronger result that Λ is contained in some tube around $\mathbb{R}^d \times Y$, where $Y = \{x \in \mathbb{R}^d : x_{d-a+1} = \dots = x_d = 0\}$. Once we have shown this, Theorem 2.11 implies the theorem.

In the case that $g_1 = g$ is compactly supported, then since E is a tube around the subspace Y and $P_{L^2(H)} M_x T_y g \neq 0$ for all $(x, y) \in \Lambda$, it follows that Λ must also be contained in a tube around $\mathbb{R}^d \times Y$.

For the general case, note that if $\{P_{L^2(H)} M_x T_y g : (x, y) \in \Lambda\}$ is a Riesz sequence, then it is in particular true that for some $\epsilon > 0$, $\inf\{\|P_{L^2(H)} M_x T_y g\| : (x, y) \in \Lambda\} > \epsilon > 0$.

Now, approximate g with a compactly supported function f so that $\|f - g\| < \epsilon$. Moreover,

$$\begin{aligned} \inf_{(x,y) \in \Lambda} \|P_{L^2(H)} M_x T_y f\| &\leq \inf_{(x,y) \in \Lambda} \|P_{L^2(H)} M_x T_y (f - g)\| + \inf_{(x,y) \in \Lambda} \|P_{L^2(H)} M_x T_y g\| \\ &\leq \epsilon + \inf_{(x,y) \in \Lambda} \|P_{L^2(H)} M_x T_y g\|. \end{aligned}$$

This implies that $\inf_{(x,y) \in \Lambda} \|P_{L^2(H)} M_x T_y g\| > 0$, which in turn implies that Λ is contained in some tube around $\mathbb{R}^d \times Y$, as desired. \square

Finally, we can give the proof of Theorem 3.3.

Proof of Theorem 3.3. (i): By a change of basis, we may assume that E is of the form given in Theorem 3.12 with $m = d - a$. Let A also be of the form given in Theorem 3.12. Then, one simply notes that $\dim_{\mathbb{R}^d}^+(\Lambda) \geq \dim_A^+(\Lambda) \geq 2d - (d - m) = d + m$, as desired.

(ii): This claim follows from Theorem 3.12 by similar arguments, we only use part (ii) instead. \square

Example 3.13. *There exists a PFFS satisfying the conditions given in Theorem 3.12 and such that $\dim_{\bar{S}}(\Lambda_k) < 2d - a$ for all $1 \leq k \leq K$.*

Proof. Let $a = 1$, $K = 1$ and $E = \mathbb{R} \times [-1/2, 1/2]$ so that $S = \mathbb{R}^3 \times \{0\} \subset \mathbb{R}^4$. We construct Λ and g such that $\mathcal{G}(G, \Lambda)$ is a tight PFFS for $L^2(E)$. Let

$$K_0 = \cup_{n=-\infty}^{\infty} [n, n+1] \times ([0, 2^{-|n|}] + 2^{2|n|})$$

Let

$$H_0 = \{0\} \times \{-2^{2|n|} + j2^{-|n|} : n \in \mathbb{N}, 0 \leq j \leq 2^{|n|} - 1\}.$$

See that $\{(K+h) \cap E : h \in H\}$ is a partition of E . It follows that setting $g = 1_K$ and $\Lambda = \mathbb{Z}^2 \times H$ yields $\mathcal{G}(G, \Lambda)$ is a PFFS for $L^2(E)$. However, $\dim_{\bar{S}}(\Lambda) = 2$. It is clear in this case that $\dim_{\mathbb{R}^{2d}}^+(\Lambda) = \infty$, as Theorem 3.12 predicts. \square

ACKNOWLEDGMENTS

The research for this paper was performed while GK was visiting the Department of Mathematics at Washington University in St. Louis, the School of Mathematics at the Georgia Institute of Technology, and NuHAG at the University of Vienna, and DS was visiting Universiteit Stellenbosch. These authors thank these departments and research groups for their hospitality and support during these visits. We are also indebted to Christopher Heil, Yang Wang, and Ed Wilson for helpful discussions.

WC was supported by European Commission grant MEIF-CT-2003-500685. GK acknowledges support from DFG research fellowship KU 1446/5 and from Preis der Justus-Liebig-Universität Gießen 2006. DS acknowledges support from the NSF through grant DMS 0354957.

REFERENCES

- [1] M. T. Barlow and S. J. Taylor, *Fractional dimension of sets in discrete spaces*, J. Phys. A: Math. Gen. **22** (1989), 2621–2626.
- [2] Y. Benyamini, *The uniform classification of Banach spaces*, Longhorn Notes, University of Texas Austin, 1984–5, 15–38.
- [3] E. J. Candès and D. L. Donoho, *New tight frames of curvelets and optimal representations of objects with piecewise C^2 singularities*, Comm. Pure Appl. Math. **57** (2004), no. 2, 219–266.
- [4] P. G. Casazza and J. C. Tremain, *The Kadison-Singer problem in mathematics and engineering*, Proc. Natl. Acad. Sci. USA **103** (2006), 2032–2039.
- [5] O. Christensen, B. Deng, and C. Heil, *Density of Gabor frames*, Appl. Comput. Harmon. Anal. **7** (1999), no. 3, 292–304.
- [6] W. Czaja, G. Kutyniok, and D. Speegle, *Geometry of sets of parameters of wave packets*, Appl. Comput. Harmon. Anal. **20** (2006), 108–125.
- [7] M. Dörfler, *Time-frequency analysis for music signals: A mathematical approach*, J. New Music Research **30** (2001), 3–12.
- [8] S. R. Forrest and T. A. Witten Jr, *Long-range correlation in smoke-particle aggregates*, J. Phys. A: Math. Gen. **12** (1979), L109–L117.
- [9] C. Heil, *History and evolution of the Density Theorem for Gabor frames*, preprint.
- [10] C. Heil, *The homogeneous approximation property of Ramanathan and Steger*, preprint.
- [11] C. Heil and G. Kutyniok, *Density of weighted wavelet frames*, J. Geom. Anal., **13** (2003), 479–493.
- [12] C. Heil and G. Kutyniok, *The Homogeneous Approximation Property for Wavelet Frames*, preprint.
- [13] G. Kutyniok, *Beurling density and shift-invariant weighted irregular Gabor systems*, Sampl. Theory Signal Image Process. **5** (2006), 163–181.
- [14] H. Landau, *Necessary density conditions for sampling and interpolation of certain entire functions*, Acta Math., **117** (1967), 37–52.
- [15] S. Li and H. Ogawa, *Pseudo-duals of frames with applications*, Appl. Comput. Harmon. Anal. **11** (2001), 289–304.
- [16] S. Li and H. Ogawa, *Pseudoframes for subspaces with applications*, J. Fourier Anal. Appl. **10** (2004), 409–431.
- [17] S. Li and H. Ogawa, *Optimal noise suppression: a geometric nature of pseudoframes for subspaces*, preprint.
- [18] J. Naudts, *Dimension of discrete fractal sets*, J. Phys. A: Math. Gen. **21** (1988), 447–452.
- [19] J. Naudts, *Reply to a paper by M. T. Barlow and S. J. Taylor*, J. Phys. A: Math. Gen. **22** (1989), 2627–2628.
- [20] J. Ramanathan and T. Steger, *Incompleteness of sparse coherent states*, Appl. Comput. Harmon. Anal., **2**, 148–153 (1995).
- [21] T. Strohmer und R. W. Heath, Jr., *Grassmannian frames with applications to coding and communication*, Appl. Comput. Harmon. Anal. **14** (2003), 257–275.
- [22] W. Sun and X. Zhou, *Density and stability of wavelet frames*, Appl. Comput. Harmon. Anal. **15** (2003), 117–133.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WROCLAW, PL. GRUNWALDZKI 2/4, 50-384 WROCLAW, POLAND, AND, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD 20742, USA

E-mail address: wojtek@math.umd.edu

INSTITUTE OF MATHEMATICS, JUSTUS-LIEBIG-UNIVERSITY GIESSEN, 35392 GIESSEN, GERMANY

E-mail address: gitta.kutyniok@math.uni-giessen.de

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, SAINT LOUIS UNIVERSITY, 221 NORTH GRAND BLVD., ST. LOUIS, MO 63103, USA

E-mail address: speegled@slu.edu